



INNOVATORS IN THE FIELD

**Foliar Fertilization**  
**Top Sidan Platinum™**

For use as sprayed foliar fertilization to supplement deficiencies

**Package: 10 liters**

**Composition:**

|                      |       |
|----------------------|-------|
| Calcium chelate (Ca) | 7.6%  |
| Boron (B)            | 0.47% |
| Silicon (Si)         | 1.0%  |

Contains: Calcium, silicon and boron in Chelate technology (PAC\*) real solution magnesium

**Recommended Dosages**

| Crop        | Dosage in cc/dunam or percentage | Application Remarks                                                                                                                                                                                                                                                                                                                                                                                                                  |
|-------------|----------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Apple, Pear | 0.3-0.5%                         | <ul style="list-style-type: none"><li>• Spray after "Pink Bulb"</li><li>• No. of applications: 6-12 according to species susceptibility to calcium deficiency.</li><li>• Duration between sprayings: 7-10 days. In young plantations with vigorous growth and twigs exceeding 45 cm, spray at least 8 times.</li><li>• The low volume in young and open plantations and the high volume in mature and tangled plantations.</li></ul> |
| Citrus      | 0.3-0.5%                         | <ul style="list-style-type: none"><li>• Spray after fruit formation</li></ul>                                                                                                                                                                                                                                                                                                                                                        |
| Pomegranate | 0.3-0.5%                         | <ul style="list-style-type: none"><li>• To prevent fruit cracking and to improve its quality, start spraying at the beginning of fruit formation. No. of applications: 4-6. Duration between applications: 14 days. When applying for the first time, combining treatment with Nanovatz Platinum is recommended in order to introduce calcium to the fruit.</li></ul>                                                                |

|                                 |          |                                                                                                                                             |                                                                                           |
|---------------------------------|----------|---------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|
| <b>Mango,<br/>Avocado</b>       | 0.3-0.5% | Spray during blooming and again, two weeks later                                                                                            |                                                                                           |
| <b>Apricot,<br/>Peach, Plum</b> | 0.3-0.5% | <ul style="list-style-type: none"> <li>• Spray from beginning of fruit formation.</li> <li>• Duration between sprayings: 14 days</li> </ul> | <ul style="list-style-type: none"> <li>• Perform 1-2 applications.</li> </ul>             |
| <b>Grapes</b>                   | 0.3-0.5% | <ul style="list-style-type: none"> <li>• Start spraying when the granule is the size of a pea.</li> <li>• Perform 4-6 sprayings.</li> </ul> | <ul style="list-style-type: none"> <li>• Duration between sprayings: 7-10 days</li> </ul> |
| <b>Tomato,<br/>Pepper</b>       | 250-400  | <ul style="list-style-type: none"> <li>• To reduce blossom end rot</li> <li>• Spray from beginning for fruit formation</li> </ul>           | <ul style="list-style-type: none"> <li>• Duration between sprayings: 7-10 days</li> </ul> |
| <b>Gourds</b>                   | 250-400  | <ul style="list-style-type: none"> <li>• Spray from beginning of fruit formation.</li> </ul>                                                | <ul style="list-style-type: none"> <li>• Duration between sprayings: 14 days</li> </ul>   |
| <b>Crucifers,<br/>Lettuce</b>   | 250-400  | <ul style="list-style-type: none"> <li>• Begin spraying when fruit begins to grow (head).</li> <li>• Perform 2-3 sprayings.</li> </ul>      | <ul style="list-style-type: none"> <li>• Duration between sprayings: 7-10 days</li> </ul> |
| <b>Celery</b>                   | 250-400  | <ul style="list-style-type: none"> <li>• Begin spraying when fruit begins to grow (head).</li> <li>• Perform 2-3 sprayings.</li> </ul>      | <ul style="list-style-type: none"> <li>• Duration between sprayings: 7 days</li> </ul>    |

\*PAC refers to Nutriag's "Poly Aldo Carbosate" Technology